arXiv:1112.0127v2 [math.CO] 7 Jul 2012

On the generalized (edge-)connectivity of graphs*

Xueliang Li, Yaping Mao, Yuefang Sun
Center for Combinatorics and LPMC-TJKLC
Nankai University, Tianjin 300071, China
lxl@nankai.edu.cn; maoyaping@ymail.com; bruceseun@gmail.com

Abstract

The generalized k-connectivity $\kappa_k(G)$ of a graph G was introduced by Chartrand et al. in 1984. It is natural to introduce the concept of generalized k-edge-connectivity $\lambda_k(G)$. For general k, the generalized k-edge-connectivity of a complete graph is obtained. For $k \geq 3$, sharp upper and lower bounds of $\kappa_k(G)$ and $\lambda_k(G)$ are given for a connected graph G of order n, that is, $1 \leq \kappa_k(G) \leq n - \lceil \frac{k}{2} \rceil$ and $1 \leq \lambda_k(G) \leq n - \lceil \frac{k}{2} \rceil$. Graphs of order n such that $\kappa_k(G) = n - \lceil \frac{k}{2} \rceil$ and $\lambda_k(G) = n - \lceil \frac{k}{2} \rceil$ are characterized, respectively. Nordhaus-Gaddum-type results for the generalized k-(edge-)connectivity are also obtained. For $k = 3$, we study the relation between the edge-connectivity and the generalized 3-edge-connectivity of a graph. Upper and lower bounds of $\lambda_3(G)$ for a graph G in terms of the edge-connectivity λ of G are obtained, that is, $\frac{3\lambda-2}{4} \leq \lambda_3(G) \leq \lambda$, and two graph classes are given showing that the upper and lower bounds are sharp. From these bounds, we obtain that $\lambda(G) - 1 \leq \lambda_3(G) \leq \lambda(G)$ if G is a connected planar graph, and the relation between the generalized 3-connectivity and generalized 3-edge-connectivity of a graph and its line graph.

Keywords: (edge-)connectivity, internally (edge-)disjoint trees, generalized (edge-)connectivity, planar graph, line graph, complementary graph.

AMS subject classification 2010: 05C40, 05C05, 05C75, 05C76.

1 Introduction

All graphs in this paper are undirected, finite and simple. We refer to book [2] for graph theoretical notation and terminology not described here.

The generalized connectivity of a graph G, which was introduced by Chartrand et al. in [4], is a natural and nice generalization of the concept of (vertex) connectivity. Let G be a nontrivial connected graph of order n and $2 \leq k \leq n$. For a set S of k vertices of G, let $\kappa(S)$ denote the maximum number ℓ of edge-disjoint trees $T_1, T_2, \cdots, T_\ell$ in G such that $V(T_i) \cap V(T_j) = S$ for every pair i, j of distinct integers with $1 \leq i, j \leq \ell$ (Note that the trees are vertex-disjoint in $G \setminus S$). A collection $\{T_1, T_2, \cdots, T_\ell\}$ of trees in G with this property is called an set of *internally disjoint trees connecting* S. The *generalized* k*-connectivity*, denoted by $\kappa_k(G)$, of G is then

---

[*] Supported by NSFC No.11071130, and “the Fundamental Research Funds for the Central Universities”

defined by $\kappa_k(G) = min\{\kappa(S) | S \subseteq V(G) \ and \ |S| = k\}$, where the minimum is taken over all k-subsets S of $V(G)$. Thus, $\kappa_2(G) = \kappa(G)$. Let $\kappa_k(G) = 0$ if G is disconnected. Results on the generalized connectivity can be found in [5, 15, 11, 12, 13, 14, 17].

A natural idea is to introduce the concept of generalized edge-connectivity. Let $\lambda(S)$ denote the maximum number ℓ of pairwise edge-disjoint trees $T_1, T_2, \cdots, T_\ell$ in G such that $V(T_i) \supseteq S$ for every $1 \leq i \leq \ell$. Then the *generalized k-edge-connectivity* $\lambda_k(G)$ of G is defined as $\lambda_k(G) = min\{\lambda(S) | S \subseteq V(G) \ and \ |S| = k\}$. Thus $\lambda_2(G) = \lambda(G)$. Let $\lambda_k(G) = 0$ if G is disconnected.

Note that the concepts of generalized connectivity and generalized edge-connectivity are different in general. The graph H in Figure 1 (a) is an example for which $\kappa_3(H) = 1$ and $\lambda_3(H) = 2$.



(a) (b)

Figure 1 (a): $\kappa_3(H) = 1, \lambda_3(H) = 2$.
Figure 1 (b): $\kappa_k(G) = 1, \lambda_k(G) = n - \lceil \frac{k}{2} \rceil$.

In addition to being natural combinatorial measures, the generalized connectivity and generalized edge-connectivity can be motivated by their interesting interpretation in practice. For example, suppose that G represents a network. If one considers to connect a pair of vertices of G, then a path is used to connect them. However, if one wants to connect a set S of vertices of G with $|S| \geq 3$, then a tree has to be used to connect them. This kind of tree with minimum order for connecting a set of vertices is usually called a Steiner tree, and popularly used in the physical design of VLSI, see [16]. Usually, one wants to consider how tough a network can be, for the connection of a set of vertices. Then, the number of totally independent ways to connect them is a measure for this purpose. The generalized k-connectivity and generalized k-edge-connectivity can serve for measuring the capability of a network G to connect any k vertices in G.

For general k, the generalized k-edge-connectivity of a complete graph is obtained. Sharp and lower bounds of $\kappa_k(G)$ and $\lambda_k(G)$ are given for a connected graph G of order n, that is, $1 \leq \kappa_k(G) \leq n - \lceil \frac{k}{2} \rceil$ and $1 \leq \lambda_k(G) \leq n - \lceil \frac{k}{2} \rceil$. We recall and use Nash-Williams's and Tutte's Theorem (for short proofs, see [10]):

Theorem 1. *(Nash-Williams [8], Tutte [9]) A multigraph G contains a system of k edge-disjoint spanning trees if and only if*

$$\|G/\mathscr{P}\| \geq k(|\mathscr{P}| - 1)$$

holds for every partition $\mathscr{P}$ of $V(G)$, where $\|G/\mathscr{P}\|$ denotes the number of edges in G between distinct blocks of $\mathscr{P}$.

By Nash-Williams's and Tutte's Theorem, graphs of order n such that $\kappa_k(G) = n - \lceil \frac{k}{2} \rceil$ and $\lambda_k(G) = n - \lceil \frac{k}{2} \rceil$ are characterized, respectively. Nordhaus-Gaddum-type results for the generalized k-connectivity and generalized k-edge-connectivity are also obtained in Section 3.

For $k = 3$, we study the relation between the edge-connectivity and the generalized 3-edge-connectivity of a graph. Kriesell in [7] showed that for any two natural numbers t, ℓ there exist smallest natural numbers $f_\ell(t)$ ($g_\ell(t)$) such that for any $f_\ell(t)$-edge-connected ($g_\ell(t)$-edge-connected) vertex set S of a graph G with $|S| \leq \ell$ ($|V(G) - S| \leq \ell$) there exists a system $\mathscr{T}$ of t edge-disjoint trees such that $S \subseteq V(T)$ for each $T \in \mathscr{T}$. He determined $f_3(t) = \lfloor \frac{8t+3}{6} \rfloor$. In Section 4, we use his this result to derive a sharp lower bound of $\lambda_3(G)$. We also give a sharp upper bound of $\lambda_3(G)$. Altogether we get that $\frac{3\lambda-2}{4} \leq \lambda_3(G) \leq \lambda$. Two graph classes are given showing that the upper and lower bounds are sharp. From these bounds, we obtain two results: one is $\lambda(G) - 1 \leq \lambda_3(G) \leq \lambda(G)$ if G is a connected planar graph, the other is the relation between the generalized 3-connectivity and generalized 3-edge-connectivity of a graph and its line graph.

2 Preliminaries

For a graph G, let $V(G)$, $E(G)$, $|G|$, $\|G\|$, $L(G)$ and $\overline{G}$ denote the set of vertices, the set of edges, the order, the size, the line graph and the complement graph of G, respectively. As usual, the *union* of two graphs G and H is the graph, denoted by $G \cup H$, with vertex set $V(G) \cup V(H)$ and edge set $E(G) \cup E(H)$. For $S \subseteq V(G)$, we denote $G \setminus S$ the subgraph by deleting the vertices of S together with the edges incident with them from G. If $S = \{v\}$, we simply write $G \setminus v$ for $G \setminus \{v\}$. If S is a subset of vertices of a graph G, the subgraph of G induced by S is denoted by $G[S]$. If M is the edge subset of G, then $G \setminus M$ denote the subgraph by deleting the edges of M. If M is a subset of edges of a graph G, the subgraph of G induced by M is denoted by $G[M]$. If $M = \{e\}$, we simply write $G \setminus e$ for $G \setminus \{e\}$. We denote by $E_G[X, Y]$ the set of edges of G with one end in X and the other end in Y. If $X = \{x\}$, we simply write $E_G[x, Y]$ for $E_G[\{x\}, Y]$.

Chartrand et al. in [5] obtained the first result in the generalized connectivity.

Theorem 2. *[5] For every two integers n and k with $2 \leq k \leq n$,*

$$\kappa_k(K_n) = n - \lceil k/2 \rceil.$$

A subtree T of G is called an *S-spanning tree*, if $S \subseteq V(T)$ holds. For distinct vertices x, y in G, let $\lambda(x, y; G)$ denote the local edge-connectivity of x and y. $S \subseteq V(G)$ is called n-edge-connected, if $\lambda(x, y; G) \geq n$ for all $x \neq y$ in S. In [7], Kriesell gave the following result:

Lemma 1. *[7] Let $t \geq 1$ be a natural number, and G be a graph, and let $\{a, b, c\} \subseteq V(G)$ be $\lfloor \frac{8t+3}{6} \rfloor$-edge-connected graph in G. Then there exists a system of t edge-disjoint $\{a, b, c\}$-spanning trees.*

It is easy to see that the definition of edge-disjoint S-spanning trees is the same as our definition of edge-disjoint trees connecting S.

The following result for planar graphs is well-known.

Lemma 2. *[2] Every planar graph G has a vertex of degree at most 5, i.e., $\delta(G) \leq 5$.*

Chartrand [6] et al. investigated the relation between the connectivity and edge-connectivity of a graph and its line graph.

Lemma 3. *[6] If G is a connected graph, then*

(1) $\kappa(L(G)) \geq \lambda(G)$, *if* $\lambda(G) \geq 2$.

(2) $\lambda(L(G)) \geq 2\lambda(G) - 2$.

(3) $\kappa(L(L(G))) \geq 2\kappa(G) - 2$.

3 Results of $\kappa_k(G)$ and $\lambda_k(G)$ for general k

After the preparation of the above section, we start to give our main results of this paper.

3.1. For complete graphs

The following two observations are easily seen.

Observation 1. *If G be a connected graph, then $\kappa_k(G) \leq \lambda_k(G) \leq \delta(G)$.*

Observation 2. *If H is a spanning subgraph of G, then $\kappa_k(H) \leq \kappa_k(G)$ and $\lambda_k(H) \leq \lambda_k(G)$.*

For general k and the complete graph K_n, $\kappa_k(K_n)$ was determined by Chartrand et al., see Theorem 2. Now we give the result for $\lambda_k(K_n)$.

Let $S = \{u_1, u_2, \cdots, u_k\} \subseteq V(G)$ and $W = V(G) \setminus S = \{w_1, w_2, \cdots, w_{n-k}\}$. Let $\mathscr{T}$ be a maximum set of edge-disjoint trees in G that connect S. Let $\mathscr{T}_1$ be the set of trees in $\mathscr{T}$ whose edges belong to $E(G[S])$, and $\mathscr{T}_2$ be the set of trees containing at least one edge of $E_G[S, W]$. Thus, $\mathscr{T} = \mathscr{T}_1 \cup \mathscr{T}_2$(Throughout this paper, $\mathscr{T}$, $\mathscr{T}_1$, $\mathscr{T}_2$ are defined as this).

Lemma 4. *Let $S \subseteq V(G)$, $|S| = k$ and T be a tree connecting S. If $T \in \mathscr{T}_1$, then T uses $k-1$ edges of $E(G[S]) \cup E_G[S, W]$; If $T \in \mathscr{T}_2$, then T uses at least k edges of $E(G[S]) \cup E_G[S, W]$.*

Proof. It is easy to see that for each T in $\mathscr{T}_1$, T uses $k-1$ edges in $E(G[S])$, namely, T uses $k-1$ edges of $E(G[S]) \cup E_G[S, W]$.

For $T \in \mathscr{T}_2$, by deleting all the vertices of T in W, we obtain some components of T in S, denoted by $C_1, C_2, \cdots, C_s$. Let $|C_i| = c_i$. Then $\|C_i\| = c_i - 1$ and $\sum_{i=1}^{s}(c_i - 1) = k - s$. Since there exists at least one edge of T between each C_i and W, where $1 \leq i \leq s$, T uses at least $(k-s) + s = k$ edges in $E(G[S]) \cup E_G[S, W]$. □

Theorem 3. *For every two integers n and k with $2 \leq k \leq n$,*

$$\lambda_k(K_n) = n - \lceil k/2 \rceil.$$

Proof. Let $G = K_n$. We choose $S \subseteq V(G)$ such that $|S| = k$. Let $|\mathscr{T}| = y$ and $|\mathscr{T}_1| = x$. From Lemma 4, each tree $T \in \mathscr{T}_1$ uses $k-1$ edges in $E(G[S]) \cup E_G[S, W]$, $|\mathscr{T}_1| = x \leq \lfloor \binom{k}{2}/(k-1) \rfloor = \lfloor \frac{k}{2} \rfloor$. Since each tree $T \in \mathscr{T}_2$ uses at least k edges in $E(G[S]) \cup E_G[S, W]$, we have $|\mathscr{T}_1|(k-1) + |\mathscr{T}_2|k \leq |E_G[S, W]| + |E(G[S])|$, that is, $x(k-1) + (y-x)k \leq \binom{k}{2} + k(n-k)$. So $\lambda_k(G) = y \leq \frac{k-1}{2} + n - k + \frac{x}{k}$.

If k is even and we assume $k = 2t$, where $t \geq 1$, then $|\mathscr{T}| = y \leq n - t - \frac{1}{2} + \frac{x}{2t} \leq n - t$ since $x \leq \lfloor \frac{k}{2} \rfloor = t$.

If k is odd and we assume $k = 2t + 1$, where $t \geq 1$, then $|\mathscr{T}| = y \leq n - t - 1 + \frac{x}{2t+1}$, which results in $y \leq n - t - 1$ since y is an integer.

From the above arguments, we conclude that $\lambda_k(K_n) \leq n - \lceil \frac{k}{2} \rceil$. Combining this with Theorem 2 and Observation 1, we have $\lambda_k(K_n) = n - \lceil \frac{k}{2} \rceil$.

□

From Theorems 2 and 3, we get that $\lambda_k(G) = \kappa_k(G)$ for a complete graph G. However, this is a very special case. Actually, $\lambda_k(G) - \kappa_k(G)$ could be very large. For example, let G be a graph obtained from two copies of the complete graph K_n by identifying one vertex in each of them, see Figure 1 (b). Then $\lambda_k(G) = n - \lceil \frac{k}{2} \rceil$, but $\kappa_k(G) = 1$.

3.2. Sharp upper and lower bounds of $\kappa_k(G)$ and $\lambda_k(G)$

Now we give the sharp bounds for $\kappa_k(G)$ and $\lambda_k(G)$:

Theorem 4. *For a connected graph G of order n and $n \geq k \geq 3$, $1 \leq \kappa_k(G) \leq n - \lceil k/2 \rceil$. Moreover, the upper and lower bounds are sharp.*

Proof. It is easy to see that $\kappa_k(G) \leq \kappa_k(K_n)$. From this together with Theorem 2, we have $\kappa_k(G) \leq n - \lceil \frac{k}{2} \rceil$. Since G is connected, then $\kappa_k(G) \geq 1$. The result holds.

One can easily check that the complete graph K_n attains the upper bound and the tree T_n attains the lower bound. □

The same upper and lower bounds can be established for the generalized k-edge-connectivity.

Theorem 5. *For a connected graph G of order n and $n \geq k \geq 3$, $1 \leq \lambda_k(G) \leq n - \lceil k/2 \rceil$. Moreover, the upper and lower bounds are sharp.*

3.3. Graphs with $\kappa_k(G) = n - \lceil k/2 \rceil$ and graphs with $\lambda_k(G) = n - \lceil k/2 \rceil$

In this section, we mainly characterize graphs with $\kappa_k(G) = n - \lceil \frac{k}{2} \rceil$ and $\lambda_k(G) = n - \lceil \frac{k}{2} \rceil$, respectively. Let us start with some lemmas, which will be used later.

Lemma 5. *For $k (n \geq k \geq 4)$ is even and $e \in E(K_n)$, $\lambda_k(K_n \setminus e) < n - \frac{k}{2}$.*

Proof. Let $G = K_n \setminus e$. We choose $S \subseteq V(G)$ such that $|S| = k$ and $K_n[S]$ containing e. Let $|\mathscr{T}| = y$ and $|\mathscr{T}_1| = x$. Since every tree $T \in \mathscr{T}_1$ uses $k-1$ edges in $E(G[S]) \cup E_G[S, W]$, $|\mathscr{T}_1| = x \leq (\binom{k}{2} - 1)/(k-1) = \frac{k}{2} - \frac{1}{k-1}$. From Lemma 4, each tree $T \in \mathscr{T}_2$ uses at least k edges of $E(G[S]) \cup E_G[S, W]$. Thus $|\mathscr{T}_1|(k-1) + |\mathscr{T}_2|k \leq |E_G[S, W]| + |E(G[S])|$, that is, $x(k-1) + (y-x)k \leq \binom{k}{2} + k(n-k) - 1$. So $\lambda_k(G) = y \leq \frac{k-1}{2} + n - k + \frac{x-1}{k} \leq n - \frac{k}{2} - \frac{1}{k-1} < n - \frac{k}{2}$. □

Lemma 6. *If $k (n \geq k \geq 3)$ is odd and M is an edge set of complete graph K_n such that $|M| \geq \frac{k+1}{2}$, $\lambda_k(K_n \setminus M) < n - \frac{k+1}{2}$.*

Proof. Let $G = K_n \setminus M$. We can choose $S \subseteq V(G)$ such that $|S| = k$ and $|M \cap \big(E(K_n[S]) \cup E_{K_n}[S,W])| \geq \frac{k+1}{2}$. Let $|\mathscr{T}| = y$ and $|\mathscr{T}_1| = x$. Since every tree $T \in \mathscr{T}_1$ uses $k-1$ edges in $E(G[S]) \cup E_G[S,W]$, $|\mathscr{T}_1| = x \leq \binom{k}{2}/(k-1) = \frac{k}{2}$. From Lemma 4, each tree $T \in \mathscr{T}_2$ uses at least k edges of $E(G[S]) \cup E_G[S,W]$. Thus $|\mathscr{T}_1|(k-1) + |\mathscr{T}_2|k \leq |E_G[S,W]| + |E(G[S])|$, that is, $x(k-1) + (y-x)k \leq \binom{k}{2} + k(n-k) - \frac{k+1}{2}$. So $\lambda_k(G) = y \leq \frac{k-1}{2} + n - k + \frac{x}{k} - \frac{k+1}{2k} \leq n - \frac{k+1}{2} - \frac{1}{2k} < n - \frac{k+1}{2}$. □

Lemma 7. *If n is odd and M is an edge set of complete graph K_n such that $0 \leq |M| \leq \frac{n-1}{2}$, then $G = K_n \setminus M$ contains $\frac{n-1}{2}$ edge-disjoint spanning trees.*

Proof. Let $\mathscr{P} = \bigcup_{i=1}^{p} V_i$ be a partition of $V(G)$ with $|V_i| = n_i (1 \leq i \leq p)$, and E be the set of edges between distinct blocks of $\mathscr{P}$ in G. The case $p = 1$ is trivial, thus we assume $p \geq 2$. Then $|E| \geq \binom{n}{2} - \sum_{i=1}^{p}\binom{n_i}{2} - |M| \geq \binom{n}{2} - \sum_{i=1}^{p}\binom{n_i}{2} - \frac{n-1}{2}$. We will show that $\binom{n}{2} - \sum_{i=1}^{p}\binom{n_i}{2} - \frac{n-1}{2} \geq \frac{n-1}{2}(p-1)$, that is, $(n-p)\frac{n-1}{2} \geq \sum_{i=1}^{p}\binom{n_i}{2}$. We only need to prove that $(n-p)\frac{n-1}{2} \geq max\{\sum_{i=1}^{p}\binom{n_i}{2}\}$. Since $f(n_1, n_2, \cdots, n_p) = \sum_{i=1}^{p}\binom{n_i}{2}$ obtain the maximum value only when $n_1 = n_2 = \cdots = n_{p-1} = 1$ and $n_p = n - p + 1$, we need inequality $(n-p)\frac{n-1}{2} \geq \binom{1}{2}(p-1) + \binom{n-p+1}{2}$, that is $(n-p)\frac{p-2}{2} \geq 0$. It is easy to see that the inequality holds. Thus, $|E| \geq \binom{n}{2} - \sum_{i=1}^{p}\binom{n_i}{2} - |M| \geq \frac{n-1}{2}(p-1)$. From Theorem 1, we know there exist $\frac{n-1}{2}$ edge-disjoint spanning trees(Note that we can use the result of Theorem 1, although Nash Williams and Tutte considered multigraphs but here we study generalized connectivity and generalized edge-connectivity for simple graphs). □

Theorem 6. *For a connected graph G of order n and $n \geq k \geq 3$, $\kappa_k(G) = n - \lceil \frac{k}{2} \rceil$ if and only if $G = K_n$ for k even; $G = K_n \setminus M$ for k odd, where M is an edge set such that $0 \leq |M| \leq \frac{k-1}{2}$.*

Proof. First we consider the case that k is even. From Theorem 2, we have $\kappa_k(K_n) = n - \frac{k}{2}$. Actually, complete graph K_n is the unique graph with this property. We only need to show that $\kappa_k(K_n \setminus e) < n - \frac{k}{2}$ for $e \in E(K_n)$. From Lemma 5 and Observation 1, we know that $\kappa_k(K_n \setminus e) \leq \lambda_k(K_n \setminus e) < n - \frac{k}{2}$ for $e \in E(K_n)$. Thus, the result holds.

Next we consider the case that k is odd.

Sufficiency: Assume that $\kappa_k(G) = n - \frac{k+1}{2}$. Since G of order n is connected, we can consider G as a graph obtained from complete graph K_n by deleting some edges. If $G = K_n \setminus M$ such that $|M| \geq \frac{k+1}{2}$, then $\kappa_k(K_n \setminus M) \leq \lambda_k(K_n \setminus M) < n - \frac{k+1}{2}$ by Observation 1 and Lemma 6, a contradiction. Thus, $G = K_n \setminus M$, where $0 \leq |M| \leq \frac{k-1}{2}$.

Necessity: We will show that $\kappa_k(G) \geq n - \frac{k+1}{2}$ if $G = K_n \setminus M$ such that $0 \leq |M| \leq \frac{k-1}{2}$. It suffices to prove that $\kappa_k(G) \geq n - \frac{k+1}{2}$ for $|M| = \frac{k-1}{2}$.

Let $S = \{u_1, u_2, \cdots, u_k\} \subseteq V(G)$ and $W = V(G) \setminus S = \{w_1, w_2, \cdots, w_{n-k}\}$. We have the following two cases to consider:

Case 1 $M \subseteq E(K_n[S]) \cup E(K_n[W])$.

Let $M' = M \cap E(K_n[S])$ and $M'' = M \cap E(K_n[W])$. Then $|M'| + |M''| = |M| = \frac{k-1}{2}$ and $0 \leq |M'|, |M''| \leq \frac{k-1}{2}$. We can consider $G[S]$ as complete graph K_k by deleting $|M'|$ edges. From Lemma 7, there exist $\frac{k-1}{2}$ edge-disjoint spanning trees in $G[S]$. Actually, these $\frac{k-1}{2}$ edge-disjoint trees are all trees connecting S in $G[S]$. All these trees together with the trees

$T_i = w_iu_1 \cup w_iu_2 \cup \cdots \cup w_iu_k (1 \leq i \leq n-k)$ form $n - \frac{k+1}{2}$ internally disjoint trees connecting S, namely, $\kappa(S) \geq n - \frac{k+1}{2}$ (Note that the trees connecting S can be edge-disjoint in $G[S]$, but must be internally disjoint in $G \setminus S$).

Case 2 $M \nsubseteq E(K_n[S]) \cup E(K_n[W])$.

In this case, there exist some edges of M in $E_{K_n}[S, W]$. Let $M' = M \cap E(K_n[S])$ and $M'' = M \cap E(K_n[W])$, and let $|M'| = m_1$ and $|M''| = m_2$. Clearly, $0 \leq m_i \leq \frac{k-3}{2} (i = 1, 2)$.

For $w_i \in W$, we let $|E_{K_n[M]}[w_i, S]| = x_i$, where $1 \leq i \leq n-k$. Without loss of generality, let $x_1 \geq x_2 \geq \cdots \geq x_{n-k}$. Actually, there exists an integer $s (1 \leq s \leq n-k)$ such that $x_{s+1} = 0$, which results in $x_{s+1} = x_{s+2} = \cdots = x_{n-k} = 0$. Thus $\sum_{i=1}^{s} x_i + m_1 + m_2 = \frac{k-1}{2}$ and $|E_G[w_i, S]| = k - x_i$. It is easy to see that $n - \frac{k+1}{2} \leq d_G(u_j) \leq n-1$ for $1 \leq j \leq k$.

Our basic idea is to seek for some edges in $G[S]$, and let them together with the edges of $E_G[S, W]$ form $n-k$ internally disjoint trees.

For $w_1 \in W$, without loss of generality, let $S = S_1 \cup S_2$ and $S_1 = \{u_1, u_2, \cdots, u_{x_1}\}$ such that $u_jw_1 \in M$ for $1 \leq j \leq x_1$. Let $S_2 = S \setminus S_1 = \{u_{x_1+1}, u_{x_1+2}, \cdots, u_k\}$. Without loss of generality, let $d_G(u_{x_1+1}) \geq d_G(u_{x_1+2}) \geq \cdots \geq d_G(u_k)$. Note that u_{x_1+1} is a vertex of maximum degree in S_2, we let $u_1' = u_{x_1+1}$. We search for the vertices of maximum degree in S_1 adjacent to $u_1' (= u_{x_1+1})$, and choose one, without loss of generality, let it be u_1. Let $e_1 = u_1'u_1$. We consider the graph $G \setminus e_1$. If $d_{G \setminus e_1}(u_{x_1+1}) \geq d_{G \setminus e_1}(u_{x_1+2})$, then we let $u_2' = u_{x_1+1}$; otherwise, we let $u_2' = u_{x_1+2}$. We search for the vertices of maximum degree in $S_1 \setminus u_1$ adjacent to u_2', and choose one, without loss of generality, let it be u_2. Let $e_2 = u_2'u_2$. Next we consider the graph $G \setminus \{e_1, e_2\}$. We choose a vertex of S_2, denoted by u_3', such that u_3' is a vertex of maximum degree in $G \setminus \{e_1, e_2\}$. We search for the vertices of maximum degree in $S_1 \setminus \{u_1, u_2\}$ adjacent to u_3', and choose one, without loss of generality, let it be u_3. Let $e_3 = u_3'u_3$. Since $d_{G[E[S_1,S_2]]}(u_j) \geq \frac{k-1}{2}$ for $1 \leq j \leq x_1$, we can proceed to find $e_4, e_5, \cdots, e_{x_1}$ by the same method. Let $M_1 = \{e_1, e_2, \cdots, e_{x_1}\}$ and $G_1 = G \setminus M_1$. It is easy to see that $d_G(u_j) - d_{G_1}(u_j) = 1 (1 \leq j \leq x_1)$ and $0 \leq d_G(u_j) - d_{G_1}(u_j) \leq x_1 (x_1 + 1 \leq j \leq k)$, which implies that $n - \frac{k+1}{2} - x_1 \leq d_{G_1}(u_j) \leq n-1$ for $1 \leq j \leq k$. The tree $T_1 = w_1u_{x_1+1} \cup w_1u_{x_1+2} \cup \cdots \cup w_1u_k \cup e_1 \cup e_2 \cup \cdots \cup e_{x_1}$ is our desired tree.

For $w_2 \in W$, without loss of generality, let $S = S_1 \cup S_2$ and $S_1 = \{u_1, u_2, \cdots, u_{x_2}\}$ such that $u_jw_2 \in M$ for $1 \leq j \leq x_2$. Let $S_2 = S \setminus S_1 = \{u_{x_2+1}, u_{x_2+2}, \cdots, u_k\}$. Without loss of generality, let u_1' be a vertex of maximum degree in S_2. We search for the vertices of maximum degree in S_1 adjacent to u_1', and choose one, without loss of generality, let it be u_1. Let $e_1 = u_1'u_1$. We consider the graph $G_1 \setminus e_1$. We choose a vertex of S_2, denoted by u_2', such that u_2' is a vertex of maximum degree in $G_1 \setminus e_1$, and proceed to search for the vertices of maximum degree in $S_1 \setminus u_1$ adjacent to u_2', and choose one, without loss of generality, let it be u_2. Let $e_2 = u_2'u_2$. Next we consider the graph $G_1 \setminus \{e_1, e_2\}$. We choose a vertex of S_2, denoted by u_3', such that u_3' is a vertex of maximum degree in $G_1 \setminus \{e_1, e_2\}$. We search for the vertices of maximum degree in $S_1 \setminus \{u_1, u_2\}$ adjacent to u_3', and choose one, without loss of generality, let it be u_3. Let $e_3 = u_3'u_3$. Since $d_{G_1[E[S_1,S_2]]}(u_j) \geq \frac{k-1}{2}$ for $1 \leq j \leq x_2$, we can proceed to find $e_4, e_5, \cdots, e_{x_2}$ by the same method. Let $M_2 = \{e_1, e_2, \cdots, e_{x_2}\}$ and $G_2 = G_1 \setminus M_2$. It is easy to see that $d_{G_1}(u_j) - d_{G_2}(u_j) = 1 (1 \leq j \leq x_2)$ and $0 \leq d_{G_1}(u_j) - d_{G_2}(u_j) \leq x_2 (x_2 + 1 \leq j \leq k)$, which implies that $n - \frac{k+1}{2} - x_1 - x_2 \leq d_{G_2}(u_j) \leq n-1$ for $1 \leq j \leq k$. The tree

$T_2 = w_2u_{x_2+1} \cup w_2u_{x_2+2} \cup \cdots \cup w_2u_k \cup e_1 \cup e_2 \cup \cdots \cup e_{x_2}$ is our desired tree. Clearly, T_2 and T_1 are internally disjoint trees connecting S.

We continue this procedure until there exists a positive integer s such that $x_{s+1} = 0$ ($|E_{K_n[M]}[w_{s+1}, S]| = 0$) for $1 \leq s \leq n-k$, we terminate this procedure and obtain s internally disjoint trees $T_1, T_2, \cdots, T_s$. The resulting graph must be $G_s = G \setminus \bigcup_{i=1}^{s} M_i$. Then $n - \frac{k+1}{2} - \sum_{i=1}^{s} x_i \leq d_{G_s}(u_j) \leq n-1$ for $1 \leq j \leq k$. Since $x_i \geq 1 (1 \leq i \leq s)$ and $n - \frac{k+1}{2} - \sum_{i=1}^{s} x_i \geq \frac{k-1}{2} - \sum_{i=1}^{s} x_i \geq 0$, the procedure can be realized. Next for $w_{s+1}, \cdots, w_{n-k}$, the trees $T_i = w_ju_1 \cup w_ju_2 \cup \cdots \cup w_ju_k (s+1 \leq j \leq n-k)$ are our desired trees.

We can consider $G_s[S] = G[S] \setminus \bigcup_{i=1}^{s} M_i$ as a graph obtained from complete graph K_k by deleting $|M'| + \bigcup_{i=1}^{s} |M_i|$ edges. Since $\sum_{i=1}^{s} x_i + m_1 + m_2 = \frac{k-1}{2}$, we have $1 \leq \bigcup_{i=1}^{s} |M_i| + m_1 \leq \frac{k-1}{2}$. From Lemma 7, there exist $\frac{k-1}{2}$ edge-disjoint trees connecting S in $G[S]$(Note that these trees can be edge-disjoint by the definition of generalized k-connectivity). These trees together with $T_1, T_2, \cdots, T_{n-k}$ form $n - \frac{k+1}{2}$ internally disjoint trees connecting S, namely, $\kappa(S) \geq n - \frac{k+1}{2}$.

From the above discussion, we get that $\kappa(S) \geq n - \frac{k+1}{2}$ for $S \subseteq V(G)$, which implies that $\kappa_k(G) \geq n - \frac{k+1}{2}$. From this together with Theorem 4, we have $\kappa_k(G) = n - \frac{k+1}{2}$.

□

Theorem 7. *For a connected graph G of order n and $n \geq k \geq 3$, $\lambda_k(G) = n - \lceil \frac{k}{2} \rceil$ if and only if $G = K_n$ for k even; $G = K_n \setminus M$ for k odd, where M is an edge set such that $0 \leq |M| \leq \frac{k-1}{2}$.*

Proof. First we consider the case that k is even. From Theorem 3 and Lemma 5, we have $\lambda_k(K_n) = n - \frac{k}{2}$ if and only if $G = K_n$.

Next we consider the case that k is odd. If $G = K_n \setminus M (0 \leq |M| \leq \frac{k-1}{2})$, then $\lambda_k(G) \geq \kappa_k(G) = n - \frac{k+1}{2}$ by Observation 1 and Theorem 6. From this together with Theorem 5, we know that $\lambda_k(G) = n - \frac{k+1}{2}$. Assume that $\lambda_k(G) = n - \frac{k+1}{2}$. Since G is connected, we can consider G as a graph obtained from complete graph K_n by deleting some edges. If $G = K_n \setminus M$ such that $|M| \geq \frac{k+1}{2}$, then $\lambda_k(G) < n - \frac{k+1}{2}$ by Lemma 6, a contradiction. So $G = K_n \setminus M$, where $0 \leq |M| \leq \frac{k-1}{2}$.

□

3.4. Nordhaus-Gaddum-type results

Alavi and Mitchem in [1] considered the Nordhaus-Gaddum-type results for connectivity and edge-connectivity. We are concerned with analogous inequalities involving the generalized k-connectivity and generalized k-edge-connectivity.

Theorem 8. *For any graph G of order n, we have $1 \leq \kappa_k(G) + \kappa_k(\overline{G}) \leq n - \lceil k/2 \rceil$. Moreover, the upper and lower bounds are sharp.*

Proof. (1) Let $\kappa_k(G) = r$ and $\kappa_k(\overline{G}) = s$. Since $G \cup \overline{G} = K_n$, $\kappa_k(K_n) \geq r + s = \kappa_k(G) + \kappa_k(\overline{G})$. This together with $\kappa_k(K_n) = n - \lceil \frac{k}{2} \rceil$ results in $\kappa_k(G) + \kappa_k(\overline{G}) \leq n - \lceil \frac{k}{2} \rceil$.

If $\kappa_k(G) + \kappa_k(\overline{G}) = 0$, then $\kappa_k(G) = \kappa_k(\overline{G}) = 0$. Thus G and $\overline{G}$ are all disconnected, which is impossible. So $\kappa_k(G) + \kappa_k(\overline{G}) \geq 1$.

We show that the bounds are sharp. The complete graph K_n attains the upper bound by Theorem 2 since $\kappa_k(K_n)+\kappa_k(\overline{K_n}) = n-\lceil\frac{k}{2}\rceil+0 = n-\lceil\frac{k}{2}\rceil$. To see that the lower bound is sharp, it suffices to take G as the complete bipartite graph $K_{1,n-1}$ since $\kappa_k(K_{1,n-1}) + \kappa_k(\overline{K_{1,n-1}}) = 1+0 = 1$. $\square$

The same Nordhaus-Gaddum-type result as that for the generalized k-connectivity can be established for the generalized k-edge-connectivity, which is stated as follows:

Theorem 9. *For any graph G of order n, we have $1 \leq \lambda_k(G) + \lambda_k(\overline{G}) \leq n - \lceil k/2\rceil$. Moreover, the upper and lower bounds are sharp.*

4 Results for $\lambda_3(G)$ and $\kappa_3(G)$

4.1. Upper and lower bounds for $\lambda_3(G)$

From now on, we focus our attention on the generalized 3-edge-connectivity. From Theorem 5, we obtain sharp upper and lower bounds of $\lambda_3(G)$, that is, $1 \leq \lambda_3(G) \leq n-2$. Now we give another sharp upper and lower bounds of $\lambda_3(G)$ by edge-connectivity, that is, $\frac{3\lambda-2}{4} \leq \lambda_3(G) \leq \lambda$, which will be used in planer graph and line graph. At first we give a sharp upper bound for $\lambda_k(G)$.

Theorem 10. *For any graph G of order n, $\lambda_k(G) \leq \lambda(G)$. Moreover, the upper bound is sharp.*

Proof. Let M be a $\lambda(G)$-edge-cut of G, where $1 \leq \lambda(G) \leq n-1$. Then $G \setminus M$ has at least two components. We can choose $S = \{v_1, v_2, \cdots, v_k\}$ so that $S \subseteq V(G)$ and two of the k vertices are in different components. Thus any tree connecting S must contain an edge in M. By the definition of $\lambda(S)$, we get $\lambda(S) \leq |M|$. So $\lambda_k(G) \leq \lambda(S) \leq |M| = \lambda(G)$.

Furthermore, we will show that the graph $G = K_k \vee (n-k)K_1 (n \geq 3k)$ satisfies $\kappa_k(G) = \lambda_k(G) = \kappa(G) = \lambda(G) = \delta(G) = k$(See Figure 2).



Figure 2: Graph G with $\kappa_k(G) = \lambda_k(G) = \kappa(G) = \lambda(G) = \delta(G) = k$.

Let $U = \overline{K_k} = \{u_1, u_2, \cdots, u_{n-k}\}$, $W = \{w_1, w_2, \cdots, w_k\}$ and S be a k-subset of vertices of G. Without loss of generality, let $|S \cap V(U)| = s (s \leq k)$. Then $|S \cap V(W)| = k-s$. Without loss of generality, let $u_i \in S (1 \leq i \leq s)$ and $w_j \in S (1 \leq j \leq k-s)$. Then the trees $T_i = w_iu_1 \cup w_iu_2 \cup \cdots \cup w_iu_s \cup u_{k+i}w_1 \cup u_{k+i}w_2 \cup \cdots \cup u_{k+i}w_{k-s} (i = 1, 2, \cdots, k-s)$ and $T_j = w_ju_1 \cup w_ju_2 \cup \cdots \cup w_ju_s \cup w_jw_1 \cup w_jw_2 \cup \cdots \cup w_jw_{k-s} (i = k-s+1, 2, \cdots, k)$ form k pairwise edge-disjoint trees connecting S, namely $\lambda(S) \geq k$. Combining this with $\lambda_k(G) \leq \lambda(G) = k$, we get $\lambda_k(G) = k$. Since the above k trees are also internally disjoint tree connecting S, we have $\kappa_k(G) = k$. So $\kappa_k(G) = \lambda_k(G) = \kappa(G) = \lambda(G) = \delta(G) = k$. Clearly, the upper bound of Theorem 10. $\square$

Next we give a sharp lower bound for $\lambda_3(G)$.

Theorem 11. *Let G be a connected graph with n vertices. For every two integers s and r with $s \geq 0$ and $r \in \{0, 1, 2, 3\}$, if $\lambda(G) = 4s + r$, then $\lambda_3(G) \geq 3s + \lceil \frac{r}{2} \rceil$. Moreover, the lower bound is sharp. We simply write $\lambda_3(G) \geq \frac{3\lambda-2}{4}$.*

Proof. Let $\lambda = \lfloor \frac{8t+3}{6} \rfloor$. From Lemma 1 we have $\lambda_3(G) \geq t$ (Note that we can use the result of Lemma 1, although Kriesell [7] considered graphs containing multiple edges but here we study generalized edge-connectivity for simple graphs).

If $\lambda = 4s$, since $\frac{8t+3}{6}$ is not an integer, then $4s < \frac{8t+3}{6}$. Thus $\lambda_3(G) \geq t > 3s - \frac{3}{8}$, which implies $\lambda_3(G) \geq 3s$. With a similar method, we can obtain that $\lambda_3(G) \geq 3s + 1$ if $\lambda = 4s + 1$, and $\lambda_3(G) \geq 3s + 2$ if $\lambda = 4s + 3$.

Note that there exists no integer t such that $4s + 2 = \lfloor \frac{8t+3}{6} \rfloor$ if $\lambda = 4s + 2$. But a graph G with $\lambda(G) = 4s + 2$ is also $(4s + 1)$-edge-connected, and so we have $\lambda_3(G) \geq 3s + 1$.

$$\lambda_3(G) \geq \begin{cases} 3s & if\ \lambda = 4s, \\ 3s+2 & if\ \lambda = 4s+3, \\ 3s+1 & if\ \lambda = 4s+1\ or\ \lambda = 4s+2. \end{cases}$$

So the result holds. Simply, we write $\lambda_3(G) \geq \frac{3\lambda-2}{4}$.

Now we give graphs attaining the lower bound.

For $\lambda = 4s$ with $s \geq 1$, we construct a graph G as follows (see Figure 3 (a)): Let $P = X_1 \cup X_2$ and $Q = Y_1 \cup Y_2$ be two cliques with $|X_1| = |Y_1| = 2s$ and $|X_2| = |Y_2| = 2s$. Let u, v be adjacent to every vertex in P, Q, respectively, and w be adjacent to every vertex in X_1 and Y_1. Finally, we finish the graph G by adding a perfect matching between X_2 and Y_2. It can be easily checked that $\lambda = 4s$.



(a) (b)

Figure 3(a): The graph with $\lambda(G) = 4s$ and $\lambda_3(G) = 3s$.
Figure 3(b): Two types of trees connecting $\{u, v, w\}$.

We consider the case $S = \{u, v, w\}$. There exist two kinds of edge-disjoint trees connecting S(see Figure 3 (b)): the tree of Type I is a path $u - v_1 - w - v_2 - v$; the tree of Type II is in T_1 or T_2, where $T_1 = uv_5 \cup v_3v_5 \cup wv_3 \cup v_5v_7 \cup v_7v$ and $T_2 = uv_6 \cup v_6v_8 \cup v_8v_4 \cup v_4w \cup v_8v$, respectively. We denote the numbers of trees of Type I and Type II by x and y, respectively. Note that $|E_G[w, X_1 \cup X_2]| = 4s$ and each tree of Type I uses two edges of $E_G[w, X_1 \cup X_2]$, we have $x \leq 2s$. Although each tree of Type II uses one edge of $E_G[w, X_1 \cup X_2]$, we have $y \leq 2s$ since each tree of Type II uses one edge of $E_G[X_2, Y_2]$ and $|E_G[X_2, Y_2]| = 2s$. Combining these with $2x + y \leq 4s$, we can derive the optimal solution $x = s$ and $y = 2s$ by solving the following integer linear programming:

$$\begin{cases} Maximize: & x + y \\ Subject\ to: & x \leq 2s,\ y \leq 2s,\ 2x + y \leq 4s, \\ and & x, y \geq 0. \end{cases}$$

Thus $\lambda(S) \geq 3s$. We can check that for other three vertices of G the number of edge-disjoint trees connecting them is not less than $3s$. So $\lambda_3(G) = 3s$ and the graph G attaining the lower bound.

For $\lambda = 4s + 1$, let $|X_1| = |Y_1| = 2s + 1$ and $|X_2| = |Y_2| = 2s$; for $\lambda = 4s + 2$, let $|X_1| = |Y_1| = 2s + 1$ and $|X_2| = |Y_2| = 2s + 1$; for $\lambda = 4s + 3$, let $|X_1| = |Y_1| = 2s + 2$ and $|X_2| = |Y_2| = 2s + 1$, where $s \geq 1$. Similarly, we can check that $\lambda_3(G) = 3s + 1$ for $\lambda = 4s + 1$; $\lambda_3(G) = 3s + 1$ for $\lambda = 4s + 2$; $\lambda_3(G) = 3s + 2$ for $\lambda = 4s + 3$.

For the case $s = 0$, we have $G = P_n$ such that $\lambda(G) = \lambda_3(G) = 1$; $G = C_n$ such that $\lambda(G) = 2$ and $\lambda_3(G) = 1$; $G = H_t$ such that $\lambda(G) = 3$ and $\lambda_3(G) = 2$, where H_t denote the graph obtained from t copies of K_4 by identifying a vertex from each of them in the way shown in Figure 4.



Figure 4: $\lambda(H_t) = 3, \lambda_3(H_t) = 2$.

□

Since $\lambda(G) \leq \delta$, by Lemma 2 we only consider a planar graph G with edge-connectivity $\lambda(G)$ at most 5. From Theorem 11, it can be deduced that for any graph (not necessarily planar) if $\lambda(G) = 1$, $\lambda_3(G) = 1$; if $\lambda(G) = 2$, $\lambda_3(G) \geq 1$; if $\lambda(G) = 3$, $\lambda_3(G) \geq 2$; if $\lambda(G) = 4$, $\lambda_3(G) \geq 3$; and if $\lambda(G) = 5$, $\lambda_3(G) \geq 4$. Therefore, the following theorem is obvious.

Theorem 12. *If G be a connected planar graph, then $\lambda(G) - 1 \leq \lambda_3(G) \leq \lambda(G)$.*

4.2. For line graph

This section investigate the relation between the generalized 3-connectivity and generalized 3-edge-connectivity of a graph and its line graph.

Theorem 13. *If G is a connected graph, then*

(1) $\lambda_3(G) \leq \kappa_3(L(G))$.

(2) $\lambda_3(L(G)) \geq \frac{3}{2}\lambda_3(G) - 2$.

(3) $\kappa_3(L(L(G)) \geq \frac{3}{2}\kappa_3(G) - 2$.

Moreover, the bounds are sharp.

Proof. For (1), let e_1, e_2, e_3 be three arbitrary distinct vertices of the line graph of G such that $\lambda_3(G) = t$ with $t \geq 1$. Let $e_1 = v_1v_1'$, $e_2 = v_2v_2'$ and $e_3 = v_3v_3'$ be those edges of G corresponding to the vertices e_1, e_2, e_3 in $L(G)$, respectively.

Consider three distinct vertices of the six end-vertices of e_1, e_2, e_3. Without loss of generality, let $S = \{v_1, v_2, v_3\}$ be three distinct vertices. Since $\lambda_3(G) = t$, there exist t edge-disjoint trees $T_1, T_2, \cdots, T_t$ connecting S in G. We define a minimal tree T connecting S as a tree connecting S whose subtree obtained by deleting any edge of T does not connect S.



Type a *Type b* *Type c*

Type d *Type e* *Type f*

Figure 5: Six types of $T_i \cup T_j$.

Choosing any two edge-disjoint minimal trees T_i and T_j $(1 \leq i, j \leq t)$ connecting S in G, we will show the trees T_i' and T_j' corresponding to T_i and T_j in $L(G)$ are internally disjoint trees. It is easy to see that $T_i \cup T_j$ has six types, as shown in Figure 5. Since T_i and T_j are edge-disjoint in G, we can find internally disjoint trees T_i' and T_j' connecting e_1, e_2, e_3 in $L(G)$. We give an example of Type c, see Figure 6. So $\kappa_3(L(G)) \geq t$ and we know the result holds.

For (2), from Theorems 10 and 11 and (2) of Lemma 3 we have $\lambda_3(L(G)) \geq \frac{3}{4}\lambda(L(G)) - \frac{1}{2} \geq \frac{3}{4}(2\lambda(G) - 2) - \frac{1}{2} = \frac{3}{2}\lambda(G) - 2 \geq \frac{3}{2}\lambda_3(G) - 2$.

For (3), from (1) and (2) of this theorem and Observation 1 we have $\kappa_3(L(L(G))) \geq \lambda_3(L(G)) \geq \frac{3}{2}\lambda_3(G) - 2 \geq \frac{3}{2}\kappa_3(G) - 2$.

One can check that the bounds are sharp if we take $G = C_n, K_6, K_6$, respectively, for (1), (2), (3). □

Let $L^0(G) = G$ and $L^1(G) = L(G)$. Then for $k \geq 2$, the *k-th* iterated line graph $L^k(G)$ is defined by $L(L^{k-1}(G))$. The next statement follows immediately from Theorem 13 and a routine application of induction.



Figure 6 (a): An example for T_i and T_j connecting S and their line graphs.
Figure 6 (b): An example for T_i' and T_j' corresponding to T_i and T_j.

Corollary 1. *If* $\kappa_3(G) = p$, *then* $\lambda_3(L^k(G)) \geq (\frac{3}{2})^k(p-\frac{8}{3})+4$ *and* $\kappa_3(L^k(G)) \geq (\frac{3}{2})^{k-1}(p-\frac{8}{3})+4$.

References

[1] Y. Alavi, J. Mitchem, *The connectivity and edge-connectivity of complementary graphs*, Lecture Notes in Math. 186(1971), 1-3.

[2] J.A. Bondy, U.S.R. Murty, *Graph Theory*, GTM 244, Springer, 2008.

[3] G. Chartrand, *A graph-theoretic approach to a communication problem*, SIAM J. Appl. Math. 14(1966), 778-781.

[4] G. Chartrand, S.F. Kappor, L. Lesniak, D.R. Lick, *Generalized connectivity in graphs*, Bull. Bombay Math. Colloq. 2(1984), 1-6.

[5] G. Chartrand, F. Okamoto, P. Zhang, *Rainbow trees in graphs and generalized connectivity*, Networks 55(4)(2010), 360-367.

[6] G. Chartrand, M. Steeart, *The connectivity of line graphs*, Math. Ann. 182(1969), 170-174.

[7] M. Kriesell, *Edge-disjoint trees containing some given vertices in a graph*, J. Combin. Theory, Ser.B, 88(2003), 53-65.

[8] C.St.J.A. Nash-Williams, *Edge-disjonint spanning trees of finite graphs*, J. London Math. Soc. 36(1961), 445-450.

[9] W. Tutte, *On the problem of decomposing a graph into n connected factors*, J. London Math. Soc. 36(1961), 221-230.

[10] S. Hakimi, J. Mitchem, E. Schmeichel, *Short proofs of theorems of Nash-Williams and Tutte*, Ars Combin. 50(1998), 257-266.

[11] S. Li, W. Li, X. Li, *The generalized connectivity of complete bipartite graphs*, Ars Combin. 104(2012).

[12] S. Li, X. Li, *Note on the hardness of generalized connectivity*, J. Combin. Optimization, in press.

[13] S. Li, X. Li, Y. Shi, *The minimal size of a graph with generalized connectivity* $\kappa_3(G) = 2$, Australasian J. Combin. 51(2011), 209-220.

[14] S. Li, X. Li, W. Zhou, *Sharp bounds for the generalized connectivity* $\kappa_3(G)$, Discrete Math. 310(2010), 2147-2163.

[15] F. Okamoto, P. Zhang, *The trees connectivity of regular complete bipartite graphs*, J. Combin. Math. Combin. Comput. 74(2010), 279-293.

[16] N.A. Sherwani, *Algorithms for VLSI physical design automation*, 3rd Edition, Kluwer Acad. Pub., London, 1999.

[17] H. Li, X. Li, Y. Sun, *The generalied* 3*-connectivity of Casesian product graphs*, Discrete Math. Theor. Comput. Sci. 14(1)(2012), 43-54.

Appendix

An example for Case 2 of Theorem 6

Let $k = 9$ and $G = K_n \setminus M$ be a connected graph of order n such that $|M| = \frac{k-1}{2} = 4$. Let $S = \{u_1, u_2, \cdots, u_9\}$, $W = G \setminus S = \{w_1, w_2, \cdots, w_{n-9}\}$ and $M = \{w_1u_2, w_1u_3, w_2u_2, u_4u_7\}$ (Figure 7 (a)). Clearly, $|E_{K_n[M]}[w_1, S]| = 2 \geq |E_{K_n[M]}[w_2, S]| = 1 > |E_{K_n[M]}[w_i, S]| = 0(3 \leq i \leq n-9)$.

For w_1, we let $S_1 = \{u_2, u_3\}$ since $w_1u_2, w_1u_3 \in M$. Let $S_2 = S \backslash S_1 = \{u_1, u_4, u_5, u_6, u_7, u_8, u_9\}$. Since $d_G(u_1) = d_G(u_5) = d_G(u_6) = d_G(u_8) = d_G(u_9) = n-1 > d_G(u_4) = d_G(u_7) = n-2$, without loss of generality, we choose $u_1' = u_1 \in S_2$. We search for the vertices adjacent to u_1' in S_1 and obtain u_2, u_3 since $u_1'u_2, u_1'u_3 \in E(G)$. Since $d_G(u_3) = n-2 > d_G(u_2) = n-3$, we choose u_3 and get $e_1 = u_1'u_3 (= u_1u_3)$. Next we consider the graph $G \setminus e_1$. Since $d_{G\setminus e_1}(u_j) = n-1 > d_{G\setminus e_1}(u_4) = d_{G\setminus e_1}(u_7) = d_{G\setminus e_1}(u_1) = n-2(j = 5, 6, 8, 9)$, without loss of generality, we choose $u_2' = u_5 \in S_2$ and proceed to search for the vertices adjacent to u_2' in $S_1 \setminus u_3$ and obtain u_2 since $S_1 \setminus u_3 = \{u_2\}$. Let $e_2 = u_2'u_2 (= u_5u_2)$. Since $|E_{K_n[M]}[w_1, S]| = 2$, we terminate this procedure. Let $M_1 = \{e_1, e_2\}$ and $G_1 = G \setminus M_1$. Thus the tree $T_1 = w_1u_1 \cup w_1u_4 \cup w_1u_5 \cup w_1u_6 \cup w_1u_7 \cup w_1u_8 \cup w_1u_9 \cup u_1u_3 \cup u_5u_2$ is our desired tree (Figure 7 (b)).



(a) (b) (c)

(d) (e)

Figure 7 Graphs for Case 2.

For w_2, we let $S_1 = \{u_2\}$ since $w_2u_2 \in M$. Let $S_2 = S \setminus S_1 = \{u_1, u_3, u_4, u_5, u_6, u_7, u_8, u_9\}$. Since $d_{G_1}(u_6) = d_{G_1}(u_8) = d_{G_1}(u_9) = n-1 > d_{G_1}(u_j) = n-2 > d_{G_1}(u_3) = n-3(j = 1, 4, 5, 7)$, without loss of generality, we choose $u_1' = u_6 \in S_2$. We search for the vertices adjacent to u_1' in S_1 and only obtain u_2. So we get $e_1 = u_1'u_2 (= u_6u_2)$. Since $|E_{K_n[M]}[w_2, S]| = 1$, we terminate this procedure. Let $M_2 = \{e_1\}$ and $G_2 = G_1 \setminus M_2$. Then the tree $T_2 = w_2u_1 \cup w_2u_3 \cup w_2u_4 \cup w_2u_5 \cup w_2u_6 \cup w_2u_7 \cup w_2u_8 \cup w_2u_9 \cup u_6u_2$ is our desired tree(Figure 7 (c)). Clearly, T_2 and T_1 are internally disjoint trees connecting S.

Since $|E_{K_n[M]}[w_j, S]| = 0$ for $3 \leq j \leq n-9$, we terminate this procedure. For $w_3, \cdots, w_{n-9}$, the trees $T_i = w_iu_1 \cup w_iu_2 \cup \cdots \cup w_iu_{n-9}(3 \leq i \leq n-9)$(Figure 7$(d)$) are our desired trees.

We can consider $G_2[S] = G[S] \setminus \{M_1, M_2\}$ as a graph obtained from complete graph K_k by deleting $1 + |M_1| + |M_2|$ edges. Since $1 + |M_1| + |M_2| = 4 = \frac{k-1}{2}$, from Lemma 7, there exist 4 edge-disjoint trees connecting S in $G[S]$ (Actually, we can give 4 edge-disjoint trees connecting S (Figure 7 (e)). For example, the trees $T_1' = u_1u_9 \cup u_2u_9 \cup u_3u_9 \cup u_3u_6 \cup u_6u_7 \cup u_6u_8 \cup u_8u_4 \cup u_5u_7$, $T_2' = u_5u_3 \cup u_5u_1 \cup u_5u_9 \cup u_9u_7 \cup u_6u_5 \cup u_3u_8 \cup u_8u_2 \cup u_4u_6$, $T_3' = u_1u_2 \cup u_2u_3 \cup u_3u_4 \cup u_4u_5 \cup u_1u_7 \cup u_7u_8 \cup u_8u_9 \cup u_9u_6$ and $T_4' = u_3u_7 \cup u_7u_2 \cup u_2u_4 \cup u_4u_1 \cup u_4u_9 \cup u_1u_6 \cup u_1u_8 \cup u_8u_5$ can be our desired trees). These trees together with $T_1, T_2, \cdots, T_{n-9}$ form $n - 5 = n - \frac{k+1}{2}$ internally disjoint trees connecting S, namely, $\kappa(S) \geq n - 5$.